




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C07C 311/16, C07D 207/09, 217/06, 295/12, 295/18, 295/22, A61K 31/18</b>		A1	(11) International Publication Number: <b>WO 99/36398</b>
			(43) International Publication Date: 22 July 1999 (22.07.99)
(21) International Application Number: PCT/GB99/00099 (22) International Filing Date: 13 January 1999 (13.01.99) (30) Priority Data: 9800750.3 14 January 1998 (14.01.98) GB (71) Applicant (for all designated States except US): ELI LILLY AND COMPANY LIMITED [GB/GB]; Kingsclere Road, Basingstoke, Hampshire RG21 6XA (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): MILUTINOVIC, Sandra [GB/GB]; 39 Devonshire Park, Reading, Berkshire RG2 7DX (GB). SIMMONDS, Robin, George [GB/GB]; 2 Plover Close, Wokingham, Berkshire RG41 3JD (GB). TUPPER, David, Edward [GB/GB]; 8 Mildenhall Close, Lower Earley, Reading, Berkshire RG6 3AT (GB). (74) Agent: HUDSON, Christopher, Mark; Eli Lilly and Company Limited, Lilly Research Centre, Erl Wood Manor, Windle- sham, Surrey GU20 6PH (GB).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published With international search report.	

(54) Title: AMINOSULPHONYLBENZAMIDE DERIVATIVES AS MODULATORS OF THE ACTIVITY OF NEURONAL CALCIUM CHANNELS



## (57) Abstract

A pharmaceutical compound of formula (I) in which the aminosulfonyl group is attached at the 3- or 4-position, and in which R<sup>1</sup> is hydrogen, C<sub>1-6</sub> Alkyl, C<sub>3-10</sub> cycloalkyl, C<sub>3-10</sub> cycloalkyl-C<sub>1-4</sub> alkyl or optionally substituted phenyl-C<sub>1-4</sub> alkyl, R<sup>2</sup> is C<sub>1-6</sub> alkyl, C<sub>3-10</sub> cycloalkyl, C<sub>3-10</sub> cycloalkyl-C<sub>1-4</sub> alkyl, optionally substituted phenyl-C<sub>1-4</sub> alkyl or -(CH<sub>2</sub>)<sub>2</sub>NR<sup>5</sup>R<sup>6</sup> where R<sup>5</sup> and R<sup>6</sup> are each hydrogen or C<sub>1-6</sub> alkyl, and R<sup>3</sup> and R<sup>4</sup> are each C<sub>1-6</sub> alkyl, C<sub>3-10</sub> cycloalkyl, C<sub>3-10</sub> cycloalkyl-C<sub>1-4</sub> alkyl, C<sub>3-6</sub> alkenyl, optionally substituted phenyl or optionally substituted phenyl-C<sub>1-4</sub> alkyl, or R<sup>1</sup> and R<sup>2</sup>, or R<sup>3</sup> and R<sup>4</sup>, or R<sup>5</sup> and R<sup>6</sup>, together with the nitrogen atom to which they are attached, form a carbocyclic group containing 4 to 7 carbon atoms optionally substituted with one to three methyl or ethyl groups and optionally containing an oxygen atom or a further nitrogen atom, said carbocyclic group being optionally fused to an optionally substituted phenyl group; or a salt thereof.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## AMINOSULPHONYLBENZAMIDE DERIVATIVES AS MODULATORS OF THE ACTIVITY OF NEURONAL CALCIUM CHANNELS


This invention relates to novel chemical compounds and their use as pharmaceuticals.

5

It is well known that chemical compounds which modulate the activity of neuronal calcium channels are potentially useful in treating disorders of the central nervous system.

10

The compounds of the invention have the following general formula:



(I)

15

in which the aminosulfonyl group is attached at the 3- or 4-position, and in which

R<sup>1</sup> is hydrogen, C<sub>1-6</sub> alkyl, C<sub>3-10</sub> cycloalkyl, C<sub>3-10</sub> cycloalkyl-C<sub>1-4</sub> alkyl or optionally substituted phenyl-C<sub>1-4</sub> alkyl,

5

R<sup>2</sup> is C<sub>1-6</sub> alkyl, C<sub>3-10</sub> cycloalkyl, C<sub>3-10</sub> cycloalkyl-C<sub>1-4</sub> alkyl, optionally substituted phenyl-C<sub>1-4</sub> alkyl or -(CH<sub>2</sub>)<sub>2</sub>NR<sup>5</sup>R<sup>6</sup> where R<sup>5</sup> and R<sup>6</sup> are each hydrogen or C<sub>1-6</sub> alkyl, and

10

R<sup>3</sup> and R<sup>4</sup> are each C<sub>1-6</sub> alkyl, C<sub>3-10</sub> cycloalkyl, C<sub>3-10</sub> cycloalkyl-C<sub>1-4</sub> alkyl, C<sub>3-6</sub> alkenyl, optionally substituted phenyl or optionally substituted phenyl-C<sub>1-4</sub> alkyl,

15

or R<sup>1</sup> and R<sup>2</sup>, or R<sup>3</sup> and R<sup>4</sup>, or R<sup>5</sup> and R<sup>6</sup>, together with the nitrogen atom to which they are attached, form a carbocyclic group containing 4 to 7 carbon atoms optionally substituted with one to three methyl or ethyl groups and optionally containing an oxygen atom or a further nitrogen atom, said carbocyclic group being optionally fused to an optionally substituted phenyl group;

20

or a salt thereof.

The compounds of the invention have been found to be active in tests that show modulation of voltage-

5 dependent calcium channels, and are thus indicated for use in the treatment of diseases in which such modulation is beneficial, in particular diseases of the central nervous system.

10 In the above formula (I), a C<sub>1-6</sub> alkyl group includes methyl, ethyl, propyl, isopropyl, butyl, tert. butyl and hexyl, and is preferably methyl or ethyl. A substituted phenyl group is phenyl substituted with one or more, for example one to three, substituents selected from, for  
15 example C<sub>1-4</sub> alkyl, especially methyl, C<sub>1-4</sub> alkoxy, especially methoxy and ethoxy, hydroxy, nitro, cyano, halo, especially chloro or fluoro, trihalomethyl, especially trifluoromethyl, carboxy and C<sub>1-4</sub> alkoxy-carbonyl. A halo atom is preferably chlorine, bromine  
20 or fluorine. A substituted phenyl group preferably has one to three substituents selected from hydroxy, C<sub>1-4</sub> alkyl, halo, nitro and trifluoromethyl. An optionally substituted phenyl-C<sub>1-4</sub> alkyl group is preferably of the formula R-(CH<sub>2</sub>)<sub>n</sub>- where R is optionally substituted  
25 phenyl and n is 1 to 4, but the linking chain can also

be branched alkylene. A C<sub>3-10</sub> cycloalkyl group is preferably, for example, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl and these groups may optionally be substituted by one or two C<sub>1-4</sub> alkyl, especially methyl, substituents. A C<sub>3-10</sub> cycloalkyl-C<sub>1-4</sub> alkyl group is one such cycloalkyl group attached to a C<sub>1-4</sub> alkyl, and is preferably of the formula R-(CH<sub>2</sub>)<sub>n</sub>- where R is cycloalkyl and n is 1 to 4. When R<sup>3</sup> or R<sup>4</sup> are C<sub>1-6</sub> alkyl they are preferably C<sub>3-6</sub> alkyl.

10 The groups R<sup>1</sup> and R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup>, and R<sup>5</sup> and R<sup>6</sup>, can form a carbocyclic ring with the nitrogen to which they are attached and optionally also contain an oxygen atom or an additional nitrogen. Preferred examples, including the nitrogen of the amino sulfonyl group, are pyrrolidino, piperazino, morpholino and especially 3,5-dimethylpiperidino.

A particular group of compounds of the invention is one of formula (I) in which R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are each C<sub>1-6</sub> alkyl, C<sub>3-10</sub> cycloalkyl, C<sub>3-10</sub> cycloalkyl-C<sub>1-4</sub> alkyl or optionally substituted phenyl-C<sub>1-4</sub> alkyl, and R<sup>1</sup> can in addition be hydrogen, or R<sup>1</sup> and R<sup>2</sup>, or R<sup>3</sup> and R<sup>4</sup>

together with the nitrogen atom to which they are attached, form a carbocyclic group as defined above.

In a preferred group of compounds  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are  
5 each  $C_{1-6}$  alkyl,  $C_{3-10}$  cycloalkyl,  $C_{3-10}$  cycloalkyl- $C_{1-4}$  alkyl or optionally substituted phenyl- $C_{1-4}$  alkyl, and  $R^1$  is in addition hydrogen.

It is preferred that  $R^1$  is hydrogen. Furthermore,  $R^3$   
10 and  $R^4$ , which can be the same or different, are preferably  $C_{1-4}$  alkyl. It is further preferred that  $R^2$  is optionally substituted phenyl- $C_{1-4}$  alkyl.

A further preferred group of compounds is one of  
15 formula (I) in which  $R^2$  is  $-(CH_2)_2NR^5R^6$ .

A further preferred group of compounds is one of  
formula (I) in which  $R^3$  or  $R^4$  is  $C_{3-6}$  alkyl or when  $R^3$   
and  $R^4$  are taken together with the nitrogen atom they  
20 form a piperidine ring which is substituted at the 3-  
and/or 5-positions with one or two methyl or ethyl  
substituents.

It will be appreciated that the compounds of the invention can contain one or more asymmetric carbon atom which gives rise to enantiomers. The compounds can be prepared as racemates or can be made from enantiomeric  
5 intermediates. Both racemates and enantiomers form part of the present invention.


It will also be understood that salts of the compounds of the invention can be prepared and such salts are  
10 included in the invention. They can be any of the well known acid addition salts. Acid addition salts are preferably the pharmaceutically acceptable non-toxic addition salts with suitable acids, such as those with inorganic acids, for example hydrochloric, hydrobromic,  
15 nitric, sulfuric or phosphoric acids, or with organic acids, such as organic carboxylic acids, for example glycollic, maleic, fumaric, malic, oxalic, tartaric, citric, salicylic or o-acetoxybenzoic acids, or organic sulfonic acids, methane sulfonic, 2-hydroxyethane  
20 sulfonic, toluene-p-sulfonic or naphthalene-2-sulfonic acids.

In addition to pharmaceutically-acceptable salts, other salts are included in the invention. They may serve as  
25 intermediates in the purification of compounds or in the preparation of other, for example pharmaceutically-



acceptable, salts, or are useful for identification, characterisation or purification.

The invention also includes a process for producing the  
5 compounds of formula (I) above which comprises reacting  
a compound of the formula:



(II)

10 where X is a leaving group such as, for example, halo or  
hydroxy, with an amine of the formula  $\text{HNR}^1\text{R}^2$ .

The reaction is preferably carried out in an organic  
solvent such as, for example, chloroform or  
15 acetonitrile, at a temperature of from  $0^\circ \text{C}$ . to  $100^\circ \text{C}$ .  
such as, for example, ambient temperature.

Intermediate compounds of formula (II) are known in the  
art and can be prepared readily by known methods. When

an acid halide is employed (X is halo such as, for example, chloro), the reaction is preferably carried out in the presence of a solid phase scavenger to absorb the acid liberated by the reaction. When the free acid is employed (X is hydroxy), a condensing reagent such as, for example, dimethylaminopropyl-ethylcarbodiimide can be employed.

Amine reactants of the formula  $\text{HNR}^1\text{R}^2$  are also well known and can be prepared readily by known methods. Those in which  $\text{R}^2$  is  $-(\text{CH}_2)_2\text{NR}^5\text{R}^6$  can, for example, be prepared by reductive amination, that is, by reacting the appropriate diamine with an aldehyde in reducing conditions.

Alternatively, such compounds in which R is  $-(\text{CH}_2)_2\text{NR}^5\text{R}^6$  can be prepared by alkylation of the corresponding compound of formula (I) in which  $\text{R}^1$  is hydrogen.

As mentioned above, the compounds of the invention are active in tests that indicate their utility in the treatment of diseases of the central nervous system. The compounds modulate the activity of calcium channels and, in particular, they block voltage sensitive calcium

channels as determined in a test based on Boot J. R.,  
et al., Specificity of autoantibodies in the Lambert-  
Eaton Myasthenic Syndrome, Ann NY Acad. Sci. (1997), in  
which measurements of calcium flux using calcium  
5 sensitive dyes are made. Compounds described in the  
following Examples were found to inhibit voltage-  
dependent calcium channels in cloned human cell lines  
expressing specific voltage-dependent calcium channels  
with an IC<sub>50</sub> of less than 10 µM.

10

The compounds of the invention are thus indicated for  
use in the treatment of anoxia, ischaemia, stroke and  
heart failure, migraine, diabetes, cognitive impairment,  
pain, epilepsy, traumatic head or spinal injury, AIDS  
15 related dementia and blindness, amnesia,  
neurodegenerative diseases such as Alzheimer's,  
Parkinson's and Huntington's diseases and age-related  
memory disorders, Down's syndrome, mood disorders, drug  
or alcohol addition withdrawal, nausea from  
20 chemotherapy, and carbon monoxide or cyanide poisoning.

The invention also includes a pharmaceutical composition  
comprising a pharmaceutically acceptable diluent or  
carrier in association with the compound of the  
25 invention or a pharmaceutically acceptable salt or ester  
thereof.

The compound may be administered by various routes, for example by the oral or rectal route, topically or parenterally, for example by injection or infusion, being usually employed in the form of a pharmaceutical composition. Such compositions are prepared in a manner well known in the pharmaceutical art and comprise at least one active compound. In making the compositions of the present invention, the active ingredient will usually be mixed with a carrier, or diluted by a carrier, and/or enclosed within a carrier which may, for example, be in the form of a capsule, sachet, paper or other container. When the carrier serves as a diluent, it may be a solid, semi-solid, or liquid material which acts as a vehicle, excipient or medium for the active ingredient. Thus, the composition may be in the form of tablets, lozenges, sachets, cachets, elixirs, suspensions, ointments containing, for example, up to 10% by weight of the compound, soft and hard gelatin capsules, suppositories, injection solutions and suspensions and sterile packaged powders.

Some examples of suitable carriers are lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, tragacanth, gelatin, syrup, methyl cellulose, methyl- and propyl-hydrobenzoate, talc magnesium stearate and mineral oil. The compositions of the injection may, as is well known

in the art, be formulated so as to provide quick, sustained or delayed release of the active ingredient after administration to the patient.

- 5 Where the compositions are formulated in unit dosage form, it is preferred that each unit dosage form contains from 5 mg to 500 mg. The term 'unit dosage form' refers to physically discrete units suitable as unit dosages for human subjects and animals, each unit  
10 containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with the required pharmaceutical carrier.

The active compound is effective over a wide dosage  
15 range and, for example, dosages per day will normally fall within the range of from 0.5 to 300 mg/kg, more usually in the range of from 5 to 100 mg/kg. However, it will be understood that the amount administered will be determined by the physician in the light of the  
20 relevant circumstances including the conditions to be treated, the choice of compound to be administered and the chosen route of administration, and therefore the above dosage ranges are not intended to limit the scope of the invention in any way.

25

The invention is illustrated by the following Preparations and Examples.

EXAMPLE 1

## 5 4-(N,N-dipropylaminosulfonyl)-N-benzyl-benzamide

To 50-100 mg polyvinylpyridine was added a 25 mM  
solution of benzylamine in chloroform (1 ml), followed  
by a 37.5 mM solution of 4-(N,N-dipropylaminosulfonyl)-  
10 benzoyl chloride in chloroform (1 ml). The mixture was  
shaken at room temperature for 4 hours.  
Aminomethylpolystyrene (100 mg, 0.1 mmole) was added and  
shaking continued for a further 16.5 hours. The mixture  
was then filtered and the resin washed with chloroform  
15 (2 x 2 ml). The combined filtrates were vacuum  
evaporated to give the required product. (TS-MS: m/z  
375, [M+H]<sup>+</sup>)

The following compounds were similarly prepared (mass  
20 spectrum values are given in brackets).

25 Thermospray Mass  
Spectrum values

---

4-(N,N-Dipropylaminosulfonyl)-N,N-dibenzyl-benzamide

(465)

	4-(N,N-Dipropylaminosulfonyl)-N-3,4-dimethoxybenzyl-benzamide	(435)
	4-(N,N-Dipropylaminosulfonyl)-N-3,5-dimethoxybenzyl-benzamide	(435)
	4-(N,N-Dipropylaminosulfonyl)-N-3-methoxybenzyl-benzamide	(405)
	4-(N,N-Dipropylaminosulfonyl)-N-3,4,5-trimethoxybenzyl-benzamide	(465)
5	4-(N,N-Dipropylaminosulfonyl)-N-4-chlorobenzyl-benzamide	(409/410)
	4-(N,N-Dipropylaminosulfonyl)-N-4-trifluoromethylbenzyl-benzamide	(443)
	4-(N,N-Dipropylaminosulfonyl)-N-4-dimethylaminobenzyl-benzamide	(418)
	4-(N,N-Dipropylaminosulfonyl)-N-4-methylbenzyl-benzamide	(389)
	4-(N,N-Dipropylaminosulfonyl)-N-3-chlorobenzyl-benzamide	(409/410)
10	4-(N,N-Dipropylaminosulfonyl)-N-3-methylbenzyl-benzamide	(389)
	4-(N,N-Dipropylaminosulfonyl)-N-3-trifluoromethylbenzyl-benzamide	(443)
	4-(N,N-Dipropylaminosulfonyl)-N-3,5-difluoromethylbenzyl-benzamide	(411)
	4-(N,N-Dipropylaminosulfonyl)-N-2,6-dimethoxybenzyl-benzamide	(435)
	4-(N,N-Dipropylaminosulfonyl)-N-2-methylbenzyl-benzamide	(389)
15	4-(N,N-Dipropylaminosulfonyl)-N-2-chlorobenzyl-benzamide	(409/410)
	4-(N,N-Dipropylaminosulfonyl)-N-2-methoxybenzyl-benzamide	(405)
	4-(N,N-Dipropylaminosulfonyl)-N-2-trifluoromethylbenzyl-benzamide	(443)
	4-(N,N-Dipropylaminosulfonyl)-N-3,4-dimethylbenzyl-benzamide	(403)
	4-(N,N-Dipropylaminosulfonyl)-N-2,6-dichlorobenzyl-benzamide	(444)
20	4-(N,N-Dipropylaminosulfonyl)-N-4-methoxyphenethyl-benzamide	(419)
	4-(N,N-Dipropylaminosulfonyl)-N-phenethyl-benzamide	(389)
	4-(N,N-Dipropylaminosulfonyl)-N-3-methoxyphenethyl-benzamide	(419)
	4-(N,N-Dipropylaminosulfonyl)-N-4-nitrophenethyl-benzamide	(434)
	4-(N,N-Dipropylaminosulfonyl)-N-2-phenylpropyl-benzamide	(403)
25	4-(N,N-Dipropylaminosulfonyl)-N-4-chlorophenethyl-benzamide	(423/424)
	4-(N,N-Dipropylaminosulfonyl)-N-4-methylphenethyl-benzamide	(403)
	4-(N,N-Dipropylaminosulfonyl)-N-2-methoxyphenethyl-benzamide	(419)
	4-(N,N-Dipropylaminosulfonyl)-N-2-chlorophenethyl-benzamide	(423/424)
	4-(N,N-Dipropylaminosulfonyl)-N-3-trifluoromethylphenethyl-benzamide	(457)
30	4-(N,N-Dipropylaminosulfonyl)-N-hexyl-benzamide	(369)
	4-(N,N-Dipropylaminosulfonyl)-N-2-methylbutyl-benzamide	(355)

	4-(4-N,N-Dipropylaminosulfonyl)benzoylmorpholine	(355)
	2-(4-N,N-Dipropylaminosulfonyl)benzoyl-6,7-dimethoxy-tetra- hydroisoquinoline	(461)
	4-(N,N-Dipropylaminosulfonyl)-N-3-methoxypropyl-benzamide	(357)
5	4-(N,N-Dipropylaminosulfonyl)-N-2-methylpropyl-benzamide	(355)
	4-(N,N-Dipropylaminosulfonyl)-N-cyclohexylmethyl-benzamide	(381)
	4-(N,N-Dipropylaminosulfonyl)-N-cyclohexyl-benzamide	(367)
	4-(N,N-Dipropylaminosulfonyl)-N-cyclopentyl-benzamide	(353)
	4-(N,N-Dipropylaminosulfonyl)-N-pentyl-benzamide	(355)
10	4-(N,N-Dipropylaminosulfonyl)-N-3-methylbutyl-benzamide	(355)
	4-(N,N-Dipropylaminosulfonyl)-N-3-phenylpropyl-benzamide	(403)
	4-(N,N-Dipropylaminosulfonyl)-N-4-tert.butylcyclohexyl-benzamide	(423)
	4-(N,N-Dipropylaminosulfonyl)-N-4-phenylbutyl-benzamide	(417)
	4-(N,N-Dipropylaminosulfonyl)-N-1-aminopropylpyrrolidine	(396)
15	4-(N,N-Dipropylaminosulfonyl)-N-3-methylcyclohexyl-benzamide	(381)
	4-(N,N-Dipropylaminosulfonyl)-N-1-benzyl-4-aminopiperidine	(458)
	4-(N,N-Dipropylaminosulfonyl)-N-cyclopropylmethyl-benzamide	(339)
	4-(N,N-Dipropylaminosulfonyl)-N-butyl-benzamide	(341)

20

EXAMPLE 2

4-(N,N-dipropylaminosulfonyl)-N-4-methoxybenzyl-  
benzamide

25

A mixture of a 75 mM solution of 4-methoxybenzylamine in  
chloroform (0.5 ml), a 75 mM solution of  
N-(dimethylaminopropyl)-N'-ethylcarbodiimide  
hydrochloride in chloroform (0.5 ml) and a 50 mM



solution of 4-(N,N-dipropylaminosulfonyl)-benzoic acid in chloroform (0.5 ml) was stirred at room temperature for 17 hours. Methanol (0.5 ml) was added with stirring and the solution applied to a methanol-washed 500 mg SCX solid phase extraction (SPE) cartridge. The cartridge was washed with methanol (4 ml) and the combined eluates vacuum evaporated to give the required product. (TS-MS: m/z 405, [M+H]<sup>+</sup>).

10

### EXAMPLE 3

4-(N,N-dibutylaminosulfonyl)-N-4-methoxybenzyl-benzamide

15 A mixture of a 200 mM solution of dibutylamine in acetonitrile (0.5 ml) and a 25 mM solution of 4-chlorosulfonylbenzoic acid in acetonitrile (1 ml) was stirred at room temperature for 18 hours. Methanol (1 ml) was then added with stirring and the solution applied to a methanol-washed 500 mg SCX SPE cartridge. The cartridge was washed with methanol (4 ml) and the combined eluates vacuum evaporated. The residue was dissolved in dichloromethane (1 ml) and a 75 mM solution of 4-methoxybenzylamine in chloroform (0.5 ml) and a 25 75 mM solution of N-(dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride in chloroform (0.5 ml)

added. This mixture was stirred at room temperature for 17 hours. Methanol (0.5 ml) was added with stirring and the solution applied to a methanol-washed 500 mg SCX SPE cartridge. The cartridge was washed with methanol  
 5 (4 ml) and the combined eluates vacuum evaporated to give the required product. (TS-MS: m/z 433, [M+H]<sup>+</sup>)

The following compounds were prepared similarly.

10

Thermospray Mass  
Spectrum values

---

	4-(N-pentylaminosulfonyl)-N-4-methoxybenzyl-benzamide	(391)
15	4-[N-(3-methylcyclohexyl)aminosulfonyl]-N-4-methoxybenzyl-benzamide	(417)
	4-[(N-butyl-N-propyl)aminosulfonyl]-N-4-methoxybenzyl-benzamide	(419)
	4-[N-(3,5-dimethylpiperidin-1-yl)aminosulfonyl]-N-4-methoxybenzyl-benzamide	(417)
	4-[(N-diisobutyl)aminosulfonyl]-N-4-methoxybenzyl-benzamide	(433)
	4-[N-(3-methylpiperidin-1-yl)aminosulfonyl]-N-4-methoxybenzyl-benzamide	(403)
20	4-[(N-methylbutyl)aminosulfonyl]-N-4-methoxybenzyl-benzamide	(405)
	4-[(4-methylpiperidin-1-yl)aminosulfonyl]-N-4-methoxybenzyl-benzamide	(403)
	4-[(3,3-dimethylpiperidin-1-yl)aminosulfonyl]-N-4-methoxybenzyl-benzamide	(417)
	4-[(N-cyclopropyl-N-propylmethyl)aminosulfonyl]-N-4-methoxybenzyl-benzamide	(417)

25

#### EXAMPLE 4

3-(N,N-dipropylaminosulfonyl)-N-3,4-dimethoxyphenethyl-benzamide

30

A mixture of a 200 mM solution of dipropylamine in acetonitrile (0.5 ml) and a 25 mM solution of 3-chlorosulfonylbenzoic acid in acetonitrile (1 ml) was stirred at room temperature for 18 hours. Methanol (1 ml) was then added with stirring and the solution applied to a methanol-washed 500 mg SCX SPE cartridge. The cartridge was washed with methanol (4 ml) and the combined eluates vacuum evaporated. The residue was dissolved in dichloromethane (1 ml) and a 75 mM solution of 3,4-dimethoxyphenethylamine in chloroform (0.5 ml) and a 75 mM solution of N-(dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride in chloroform (0.5 ml) added. This mixture was stirred at room temperature for 17 hours. Methanol (0.5 ml) was added with stirring and the solution applied to a methanol-washed 500 mg SCX SPE cartridge. The cartridge was washed with methanol (4 ml) and the combined eluates vacuum evaporated to give the required product. (TS-MS: m/z 449, [M+H]<sup>+</sup>)

20

EXAMPLE 5

(1) 4-[(N,N-di-n-propylamino)sulfonyl]-benzoic acid

25 To a stirred solution of di-n-propylamine (3.03 g, 0.03 mole) in dry tetrahydrofuran (20 ml) at 0° C.

(ice/salt bath), was added 4-chlorosulfonylbenzoic acid (2.2 g, 0.01 mole). Stirring was continued for 1 hour. Ice water was added cautiously and the reaction made acid with 2NHCl. The 4-[(N,N-di-n-propylamino)sulfonyl]-benzoic acid was collected by filtration as a white solid which was dried *in vacuo* at 40° C.

Similarly prepared were:

10

- 3-[(N,N-di-n-propylamino)sulfonyl]-benzoic acid
- 4-[(N-phenyl-N-n-propylamino)sulfonyl]-benzoic acid
- 4-[(N-phenyl-N-n-allylamino)sulfonyl]-benzoic acid
- 3-[N-(3,3-dimethylpiperidin-1-yl)sulfonyl]-benzoic acid
- 15 4-[N-(3,3-dimethylpiperidin-1-yl)sulfonyl]-benzoic acid
- 4-[(N-phenyl-N-n-butylamino)sulfonyl]-benzoic acid
- 3-[(N-phenyl-N-n-propylamino)sulfonyl]-benzoic acid
- 3-[N-(3-ethylpiperidin-1-yl)sulfonyl]-benzoic acid
- 3-[(N-phenyl-N-methyl)sulfonyl]-benzoic acid
- 20 4-[N-(3-methylpiperidin-1-yl)sulfonyl]-benzoic acid

(2) 4-[(N-di-n-propylamino)sulfonyl]-N-4-methoxybenzyl-benzamide

25

To a solution of 4-[(N,N-di-n-propylamino)sulfonyl]-benzoic acid (2.85 g, 0.01 mole) in dry dichloromethane (ml) at 0° C. was added oxalyl chloride (2.54 g, 0.02 mole) and dimethylformamide (4 drops). The reaction mixture was stirred for 2 hours. The reaction was evaporated to dryness *in vacuo*. The resulting acid chloride was added to a stirred solution of *p*-methoxybenzylamine (1.51 g, 0.011 mole) and triethylamine (1.11 g, 0.011 mole) in dry tetrahydrofuran (25 ml) at 0-5° C. After stirring for 4 hours the reaction was poured into ice water and extracted with ethyl acetate. The solvent was washed with brine, dried and evaporated to dryness *in vacuo*. Chromatography on flash silica using 10% ethyl acetate/dichloromethane gave 4-[(N,N-di-n-propylamino)sulfonyl]-N-4-methoxybenzyl-benzamide as a white solid. M.p. 132-134° C.

Similarly prepared were:

20

3-[(N,N-di-n-propylamino)sulfonyl]-N-4-methoxybenzyl-benzamide. M.p. 132-134° C.

4-[(N-phenyl-N-n-propylamino)sulfonyl]-N-4-methoxybenzyl-benzamide. M.p. 112-114° C.

25

4-[(N-phenyl-N-n-butylamino)sulfonyl]-N-n-hexyl-benzamide. M.p. 84-86° C.

- 4-[(N-phenyl-N-n-allylamino)sulfonyl]-N-n-hexyl-benzamide. M.p. 90-92° C.
- 4-[(N-phenyl-N-n-propylamino)sulfonyl]-N-n-hexyl-benzamide. M.p. 92-94° C.
- 5 3-[N-(3,3-dimethylpiperidin-1-yl)sulfonyl]-N-4-fluorobenzyl-benzamide. M.p. 125° C.
- 4-[N-(3,3-dimethylpiperidin-1-yl)sulfonyl]-N-4-fluorobenzyl-benzamide. M.p. 138-140° C.
- 3-[(N,N-di-n-propylamino)sulfonyl]-N-4-fluorobenzyl-  
10 benzamide. M.p. 84-86° C.
- N-methyl-3-[N-(3,3-dimethylpiperidin-1-yl)sulfonyl]-N-4-fluorobenzyl-benzamide. M.p. <50° C.
- N-benzyl-3-[N-(3,3-dimethylpiperidin-1-yl)sulfonyl]-N-4-fluorobenzyl-benzamide. M.p. 112° C.
- 15 4-[(N-phenyl-N-n-butylamino)sulfonyl]-N-4-fluorobenzyl-benzamide. M.p. 128-130° C.
- 3-[(N-phenyl-N-n-propylamino)sulfonyl]-N-4-fluorobenzyl-benzamide. M.p. 99° C.
- 4-[(3,3-dimethylpiperidin-1-yl)sulfonyl]-N-(2-[(4-fluorophenyl)methyl] amino)ethyl)benzamide. Maleate.  
20 M.p. 132-134° C.
- 3-[N-(3-ethylpiperidin-1-yl)sulfonyl]-N-4-fluorobenzyl-benzamide. 405 (M+H)<sup>+</sup>
- N-[2-(dimethylamino)ethyl]-3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]-N-4-fluorobenzyl-benzamide maleate. M.p.  
25 126° C.

- N*-[2-(dimethylamino)ethyl]-3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]-*N*-cyclohexylmethyl-benzamide. M.p. 122° C.
- N*-[2-(dimethylamino)ethyl]-4-[(*N*-phenyl-*N*-n-propylamino)sulfonyl]-*N*-cyclohexylmethyl-benzamide
- 5 maleate. M.p. 140-142° C.
- N*-[2-(pyrrolidino)ethyl]-3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]-*N*-isoamyl-benzamide hydrochloride. M.p. 179° C.
- N*-[2-(pyrrolidino)ethyl]-4-[(3-ethylpiperidin-1-yl)sulfonyl]-*N*-isoamyl-benzamide malea. M.p. 166-168° C.
- 10 *N*-[3-(pyrrolidino)propyl]-3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]-*N*-isoamyl-benzamide hydrochloride. M.p. 155° C.
- 15 *N*-[3-(pyrrolidino)propyl]-3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]-*N*-cyclohexylmethyl-benzamide. M.p. 124° C.
- N*-[2-(*N*-methyl-pyrrolidin-2-yl)ethyl]-3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]-*N*-isoamyl-benzamide maleate. M.p. 115-117° C.
- 20 *N*-[2-(piperidin-1-yl)ethyl]-3-[(3-methylpiperidin-1-yl)sulfonyl]-*N*-(2-4-methoxyphenethyl)-benzamide. M.p. 210° C.
- N*-[2-(piperidin-1-yl)ethyl]-3-[(3-methylpiperidin-1-yl)sulfonyl]-*N*-(2-4-methoxyphenethyl)-benzamide
- 25 hydrochloride. M.p. 205° C.

The following Examples illustrate typical formulations containing a compound of the invention.

5

EXAMPLE 6

Tablets each containing 10 mg of active ingredient are made up as follows:

10

Active ingredient	10 mg
Starch	160 mg
Microcrystalline cellulose	100 mg
Polyvinylpyrrolidone (as 10% solution in water)	13 mg
15 Sodium carboxymethyl starch	14 mg
Magnesium stearate	3 mg
	<hr/>
Total	300 mg
	<hr/>

20

The active ingredient, starch and cellulose are mixed thoroughly. The solution of polyvinylpyrrolidone is mixed with the resultant powders and passed through a sieve. The granules so produced are dried and re-passed  
25 through a sieve. The sodium carboxymethyl starch and magnesium stearate are then added to the granules which,



after mixing, are compressed on a tablet machine to yield tablets each weighing 300 mg.

EXAMPLE 7

5

Capsules each containing 20 mg of active ingredient are made as follows:

	Active ingredient	20 mg
10	Dried starch	178 mg
	Magnesium stearate	2 mg
		_____
	Total	200 mg
		_____

15

The active ingredient, starch and magnesium stearate are passed through a sieve and filled into hard gelatine capsules in 200 mg quantities.

20

EXAMPLE 8

Capsules each containing 20 mg of medicament are made as follows:

25

	Active ingredient	20 mg
	Lactose	171 mg
	Sodium lauryl sulphate	2 mg
	Sodium starch glycollate	6 mg
5	Magnesium stearate	1 mg

---

200 mg

---

- 10 The active ingredient, lactose, sodium lauryl sulphate and sodium starch glycollate are mixed thoroughly. The blend is mixed with the magnesium stearate and filled into hard gelatine capsules in 200 mg quantities.

15 EXAMPLE 9

Tablets each containing 20 mg and medicaments are made as follows:

20	Active ingredient	20 mg
	Lactose	103 mg
	Microcrystalline cellulose	150 mg
	Hydroxypropylmethylcellulose	15 mg
	Sodium starch glycollate	9 mg
25	Magnesium stearate	3 mg

---


300 mg

---

- 5 The active ingredient, lactose, microcrystalline cellulose, sodium starch glycollate and hydroxypropylmethylcellulose are passed through a sieve and blended together. Water is added to the blended powders to form a damp mass. The damp mass is passed
- 10 through a coarse screen, dried, then re-screened. The dried granules are mixed with the magnesium stearate and compressed into tablets of 300 mg weight.

CLAIMS

1. A compound of the formula



(I)

5

in which the aminosulfonyl group is attached at the  
3- or 4-position, and in which

- 10  $R^1$  is hydrogen,  $C_{1-6}$  alkyl,  $C_{3-10}$  cycloalkyl,  $C_{3-10}$   
cycloalkyl- $C_{1-4}$  alkyl or optionally substituted  
phenyl- $C_{1-4}$  alkyl,

- $R^2$  is  $C_{1-6}$  alkyl,  $C_{3-10}$  cycloalkyl,  $C_{3-10}$   
15 cycloalkyl- $C_{1-4}$  alkyl, optionally substituted  
phenyl- $C_{1-4}$  alkyl or  $-(CH_2)_2NR^5R^6$  where  $R^5$  and  $R^6$   
are each hydrogen or  $C_{1-6}$  alkyl, and

R<sup>3</sup> and R<sup>4</sup> are each C<sub>1-6</sub> alkyl, C<sub>3-10</sub> cycloalkyl,  
C<sub>3-10</sub> cycloalkyl-C<sub>1-4</sub> alkyl, C<sub>3-6</sub> alkenyl,  
optionally substituted phenyl or optionally  
5 substituted phenyl-C<sub>1-4</sub> alkyl,

or R<sup>1</sup> and R<sup>2</sup>, or R<sup>3</sup> and R<sup>4</sup>, or R<sup>5</sup> and R<sup>6</sup>, together  
with the nitrogen atom to which they are attached,  
form a carbocyclic group containing 4 to 7 carbon  
10 atoms optionally substituted with one to three  
methyl or ethyl groups and optionally containing an  
oxygen atom or a further nitrogen atom, said  
carbocyclic group being optionally fused to an  
optionally substituted phenyl group;

15


or a salt thereof.

2. A compound according to Claim 1 in which R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>  
and R<sup>4</sup> are each C<sub>1-6</sub> alkyl, C<sub>3-10</sub> cycloalkyl, C<sub>3-10</sub>  
20 cycloalkyl-C<sub>1-4</sub> alkyl or optionally substituted  
phenyl-C<sub>1-4</sub> alkyl, and R<sup>1</sup> can in addition be  
hydrogen, or R<sup>1</sup> and R<sup>2</sup>, or R<sup>3</sup> and R<sup>4</sup> together with  
the nitrogen atom to which they are attached, form  
a carbocyclic group.

3. A compound according to Claim 2 in which  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are each  $C_{1-6}$  alkyl,  $C_{3-10}$  cycloalkyl,  $C_{3-10}$  cycloalkyl- $C_{1-4}$  alkyl or optionally substituted phenyl- $C_{1-4}$  alkyl, and  $R^1$  can in addition be hydrogen.
4. A compound according to Claim 3 in which  $R^1$  is hydrogen,  $R^2$  is optionally substituted phenyl- $C_{1-4}$  alkyl and  $R^3$  and  $R^4$  are  $C_{1-6}$  alkyl.
5. A compound according to Claim 1 in which  $R^2$  is  $-(CH_2)_2NR^5R^6$ .
6. A compound according to Claim 1 or 5 in which  $R^3$  or  $R^4$  is  $C_{3-6}$  alkyl or when  $R^3$  and  $R^4$  are taken together with the nitrogen atom they form a piperidine ring which is substituted at the 3- and/or 5-positions with one or two methyl or ethyl substituents.
7. A pharmaceutical formulation comprising a compound according to any of Claims 1 to 6 or a

pharmaceutically acceptable salt thereof, together with a diluent or carrier therefor.

8. A compound according to any of Claims 1 to 6, for  
5 use as a pharmaceutical.
9. Use of a compound according to any of Claims 1 to 6, in the manufacture of a medicament for treating a disease of the central nervous system.
- 10
10. A process for producing a compound according to Claim 1, which comprises reacting a compound of the formula



(II)

15 where X is a leaving group, with an amine of the formula HNR<sup>1</sup>R<sup>2</sup>.

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07C311/16 C07D207/09 C07D217/06 C07D295/12 C07D295/18  
C07D295/22 A61K31/18

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07C C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	C.H ANDREWS, ET AL.: "Experimental chemotherapy of typhus. Antirickettsial action of p-sulphamylbenzamidine and related compounds" PROCEEDINGS OF THE ROYAL SOCIETY OF LONDON, SERIES B - BIOLOGICAL SCIENCES, vol. 133, 1946, pages 20-60, XP002100404 London, GB see page 47, line 12 - line 22 ---	1-3
X	P. BEAK, ET AL.: "The tertiary amide as an effective director of ortho lithiation" JOURNAL OF ORGANIC CHEMISTRY, vol. 47, no. 1, 1 January 1982, pages 34-46, XP002100405 Washington, DC, US see compound 22 --- -/--	1-3

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

## \* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

20 April 1999

Date of mailing of the international search report

06/05/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

English, R



## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	K. YONEYAMA, ET AL.: "Phytotoxic activity of N-benzylbenzenesulphonamides" AGRICULTURAL AND BIOLOGICAL CHEMISTRY, vol. 48, no. 2, February 1984, pages 491-498, XP002100406 New York, US see page 493, lines 29-34 and compound II where R <sub>1</sub> , Y = H, R <sub>2</sub> = Me, X = CONMe <sub>2</sub> -----	1-3
A	US 5 591 754 A (H.-J. LANG, ET AL.) 7 January 1997 see column 1 - column 3 -----	1-10

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5591754 A	07-01-1997	AT 147375 T	15-01-1997
		AU 661611 B	27-07-1995
		AU 4746093 A	31-03-1994
		CA 2106613 A	23-03-1994
		CN 1087900 A	15-06-1994
		DE 59305042 D	20-02-1997
		DK 589336 T	16-06-1997
		EP 0589336 A	30-03-1994
		ES 2097409 T	01-04-1997
		FI 934109 A	23-03-1994
		GR 3022549 T	31-05-1997
		HU 70186 A	28-09-1995
		JP 6228082 A	16-08-1994
		NO 933351 A,B,	23-03-1994
		NZ 248703 A	21-12-1995
		ZA 9306960 A	15-04-1994

---

BEST AVAILABLE COPY